

1) (2 points) Solve for the variable in $x^4 - 1 = 0$. *Hint, there are four answers.*

$$(x^2 - 1)(x^2 + 1) = 0$$

$$x^2 - 1 = 0 \quad x^2 + 1 = 0$$

$$x = \pm 1 \quad x^2 = -1 \quad x = \pm i$$

2) (2 points each) For the function $f(x) = 2x^2 - 8x + 1$, determine...

a) If it opens up or down. How do you know?

up $a = 2 > 0$

b) The coordinates of the vertex:

$$x = -\frac{b}{2a} = -\frac{(-8)}{2(2)} = 2$$

$$f(2) = 2(2)^2 - 8(2) + 1 = -7$$

$(2, -7)$

c) The domain:

\mathbb{R}

d) The range:

$[-7, \infty)$

e) Interval of increase:

$(2, \infty)$


f) Interval of decrease:

$(-\infty, 2)$

3) (5 points) Dolores Umbridge is upset with the new crop of first-year students at Hogwarts. She decides to enclose them in an evil rectangular garden of doom near the Whomping Willow. She wants to create 3 adjacent pens that which are enclosed on all sides using 200 feet of fence. What should the dimensions of the enclosure be to maximize area? Also, what is the maximum area? Be sure to draw a picture for this scenario.



Ms. Umbridge sips her tea whilst gleefully judging her students.



$$4x + 2y = 200 \Rightarrow y = 100 - 2x$$

$$A = xy = x(100 - 2x) = -2x^2 + 100x$$

$$x = -\frac{b}{2a} = -\frac{100}{2(-2)} = 25 \text{ ft}$$

$$y = 100 - 2(25) = 50 \text{ ft}$$

$$A(25) = 25 \cdot 50 = 1250 \text{ ft}^2$$

or

$$-2(25)^2 + 100(25)$$

4) (2 points each) Solve for the variable. $\frac{3x-7}{x^2-9} - \frac{5x^2-7}{x-3} = \frac{7}{x+3} \cdot x^2-9$

$$3x-7 - 5(x+3) = 7(x-3) \Rightarrow x = 1/9$$


$$3x-7 - 5x-15 = 7x-21$$

$$-2x - 22 = 7x - 21$$

21


5) (2 points each) Give an example of a function which will have similar arrowheads to the function below:

a)



$f(x) = \underline{\text{neg } x^{\text{even}}}$

b)



$f(x) = \underline{\text{pos } x^{\text{odd}}}$


leading term

6) (3 points each) For the function $f(x) = (x+4)(x-2)^2(x-5)\dots$

a) Find the leading term and state which quadrants the arrowheads will be in:

$y = x^4 \rightarrow \text{even}$
 $\hookrightarrow \text{pos} \rightarrow \text{Q I \& II}$

c) Sketch the graph based on parts a and b:



b) Fill in the chart:

Zero	Multiplicity	Touch/Cross
-4	1	C
2	2	T
5	1	C


$\hookrightarrow -3x^2(x^2-4)$

7) (3 points each) For the function $f(x) = -3x^4 + 12x^2 \dots = -3x^2(x+2)(x-2)$

a) Find the leading term and state which quadrants the arrowheads will be in:

$\text{neg } x^{\text{even}} \rightarrow \text{Q III \& IV}$

c) Sketch the graph based on parts a and b:



b) Fill in the chart:

Zero	Multiplicity	Touch/Cross
0	2	T
-2	1	C
2	1	C

22

8) (2 points each) Form a polynomial function of degree four that meets the following requirements. **Be sure to leave your answer in factored form:**

a) Has zeros at 8, 3, 5, and 0:

$$f(x) = (x-8)(x-3)(x-5)(x-0)$$

or just x

b) Has the same zeros and multiplicity as in part a but is a different function:

$$g(x) = \underline{5.36} (x-8)(x-3)(x-5)(x-0)$$

any nonzero #

c) Has a zero at $2+3i$, and 8 is a zero of multiplicity 2:

$$h(x) = (x - (2+3i))(x - (2-3i))(x-8)^2$$

9) (3 pts a; 2 pts others) Consider the functions $f(x) = 6x^3 + x^2 - 12x + 5$ and

$$g(x) = 3x^2 + 2x - 5.$$

a) Divide $f(x)$ by $g(x)$ using long division:

$$\begin{array}{r} 2x - 1 \\ 3x^2 + 2x - 5 \overline{) 6x^3 + x^2 - 12x + 5} \\ \underline{-(6x^3 + 4x^2 - 10x)} \\ -3x^2 - 2x + 5 \\ \underline{-(-3x^2 - 2x + 5)} \\ 0 \end{array}$$

b) Based on your work in part a, was $g(x)$ a factor of $f(x)$? Why or why not?

Yes
Remainder was zero!

c) What is the equation of the oblique asymptote of the rational function $y = \frac{6x^3 + x^2 - 12x + 5}{3x^2 + 2x - 5}$?

$$y = 2x - 1$$

13

10) (7 points each) Factor the polynomial completely by first listing the possible rational roots and then using synthetic division and your calculator.

a) $f(x) = x^3 - 5x^2 - 12x + 36$

$\frac{p}{q} = \pm 1, 2, 3, 4, 6, 9, 12, 18, 36$

$$\begin{array}{r} -3 \overline{) 1 \quad -5 \quad -12 \quad 36} \\ \underline{-3 \quad 24 \quad -36} \\ 1 \quad -8 \quad 12 \quad 0 \\ 2 \overline{) 1 \quad -8 \quad 12 \quad 0} \\ \underline{2 \quad -12} \\ 1 \quad -6 \quad 0 \end{array}$$

$f(x) = (x+3)(x-2)(x-6)$

b) $g(x) = x^4 - 2x^3 - 10x^2 + 16x + 40$

$\frac{p}{q} = \pm 1, 2, 4, 5, 8, 10, 20, 40$

$$\begin{array}{r} -2 \overline{) 1 \quad -2 \quad -10 \quad 16 \quad 40} \\ \underline{-2 \quad 8 \quad 4 \quad -40} \\ -2 \overline{) 1 \quad -4 \quad -6 \quad 20 \quad 0} \\ \underline{-2 \quad 12 \quad -20} \\ 1 \quad -6 \quad 10 \quad 0 \end{array}$$

$= (x+2)^2(x^2 - 6x + 10)$ quadratic formula.

$g(x) = (x+2)^2(x - (3+i))(x - (3-i))$

11) (3 points each) For the function $f(x) = \frac{x-2}{x^2-x-2}$, find...

a) The domain:

$x^2 - x - 2 = 0$
 $(x-2)(x+1) = 0$
 $x \neq 2, -1$

b) The x- and y-intercepts:

$\frac{x\text{-int}}{x-2=0}$
 $x=2$
 \emptyset
 not in domain

$\frac{y\text{-int}}{f(0) = \frac{-2}{-2}}$
 $= 1$
 $(0, 1)$

c) Any vertical asymptotes and holes:

$\frac{x=-1}{-1-2 \neq 0}$
 VA
 $x=-1$


$\frac{x=2}{2-2=0 \checkmark}$
 $\frac{x-2}{(x-2)(x+1)} = \frac{1}{x+1}$
 @ $x=2$
 $\frac{1}{2+1} = \frac{1}{3}$
 Ho: $(2, 1/3)$

d) Any horizontal or oblique asymptotes:

HA
 $y=0$

e) Sketch a graph based on the above.

Hint: you should see a transformation of $y = \frac{1}{x}$ in your work. Use that!



Handwritten signature

12) (2 points each) Fill in the blank:

a) If c is a zero of a function f , then $f(c) =$ bacon, and pancake is a factor.

b) Numbers not in the domain of a rational function lead to making bacon pancakes.

13) (3 points each) Short answer. Clearly explain how to find the following algebraically:

a) Vertical Asymptotes and Holes:

b) Horizontal and Oblique Asymptotes:

take some
bacon


I'll put it
in a pancake

14) (3 points each) Solve for the variable. Write answer in interval notation:

$$(x+4)(x-2)^2 \leq 0$$

$$(x+4)(x-2)^2 = 0$$

$$x = -4 \quad x = 2$$



$$(-\infty, -4] \cup \{2\}$$

evil is

Extra Credit (2 points):

Find the equation of a rational function in **factored form** that has the following properties:

a) Hole at $x = 9$

b) Vertical Asymptotes at $x = -3$ and $x = 5$

c) x -intercepts at $x = \frac{1}{7}$ and $x = -6$

d) Horizontal asymptote at $y = 7$

Hmmm

15/17